

VOLUME 12 PART 1

DECEMBER 1976

ISSN 0303-2515

QH
1
S67X
NH



ANNALS

OF THE SOUTH AFRICAN
MUSEUM

CAPE TOWN



INSTRUCTIONS TO AUTHORS

1. MATERIAL should be original and not published elsewhere, in whole or in part. When accepted, copyright becomes the property of the Trustees of the South African Museum.

2. LAYOUT should be as follows:

- (a) *Centred masthead to consist of*
Title: informative but concise, without abbreviations and not including the names of new genera or species
Author's(s') name(s)
Address(es) of author(s) (institution where work was carried out)
Number of illustrations (figures, enumerated maps and tables, in this order)
- (b) *Abstract* of not more than 200 words, intelligible to the reader without reference to the text
- (c) *Table of contents* giving hierarchy of headings and subheadings
- (d) *Introduction*
- (e) *Subject-matter* of the paper, divided into sections to correspond with those given in table of contents
- (f) *Summary*, if paper is lengthy
- (g) *Acknowledgements*
- (h) *References*
- (i) *Abbreviations*, where these are numerous

3. MANUSCRIPT, to be submitted in triplicate, should be typewritten and neat, double spaced with 2,5 cm margins all round. First lines of paragraphs should be indented. Tables and a list of legends for illustrations should be typed separately, their positions indicated in the text. All pages should be numbered consecutively.

Major headings of the paper are centred capitals; first subheadings are shouldered small capitals; second subheadings are shouldered italics; third subheadings are indented, shouldered italics. Further subdivisions should be avoided, as also enumeration (never roman numerals) of headings and abbreviations.

Footnotes should be avoided unless they are short and essential.

Only generic and specific names should be underlined to indicate italics; all other marking up should be left to editor and publisher.

4. ILLUSTRATIONS should be reducible to a size not exceeding 12 × 18 cm (19 cm including legend); the reduction or enlargement required should be indicated; originals larger than 35 × 47 cm should not be submitted; photographs should be rectangular in shape and final size. A metric scale should appear with all illustrations, otherwise magnification or reduction should be given in the legend; if the latter, then the final reduction or enlargement should be taken into consideration.

All illustrations, whether line drawings or photographs, should be termed figures (plates are not printed; half-tones will appear in their proper place in the text) and numbered in a single series. Items of composite figures should be designated by capital letters; lettering of figures is not set in type and should be in lower-case letters.

The number of the figure should be lightly marked in pencil on the back of each illustration.

5. REFERENCES cited in text and synonymies should all be included in the list at the end of the paper, using the Harvard System (*ibid.*, *idem*, *loc. cit.*, *op. cit.* are not acceptable):

(a) Author's name and year of publication given in text, e.g.:

- 'Smith (1969) describes ...'
'Smith (1969: 36, fig. 16) describes ...'
'As described (Smith 1969a, 1969b; Jones 1971)
'As described (Haughton & Broom 1927) ...'
'As described (Haughton *et al.* 1927) ...'

Note: no comma separating name and year
pagination indicated by colon, not p.
names of joint authors connected by ampersand
et al. in text for more than two joint authors, but names of all authors given in list of references.

(b) Full references at the end of the paper, arranged alphabetically by names, chronologically within each name, with suffixes *a*, *b*, etc. to the year for more than one paper by the same author in that year, e.g. Smith (1969a, 1969b) and not Smith (1969 1969a).

For books give title in italics, edition, volume number, place of publication, publisher.

For journal article give title of article, title of journal in italics (abbreviated according to the *World list of scientific periodicals*, 4th ed. London: Butterworths, 1963), series in parentheses, volume number, part number (only if independently paged) in parentheses, pagination (first and last pages of article).

Examples (note capitalization and punctuation)

- BULLOUGH, W. S. 1960. *Practical invertebrate anatomy*. 2nd ed. London: Macmillan.
FISCHER, P.-H. 1948. Données sur la résistance et de la vitalité des mollusques. — *J. Conch.*, Paris 88: 100-140.
FISCHER, P.-H., DUVAL, M. & RAFFY, A. 1933. Etudes sur les échanges respiratoires des littorines. — *Archs Zool. exp. gen.* 74: 627-634.
KOHN, A. J. 1960a. Ecological notes on *Conus* (Mollusca: Gastropoda) in the Trincomalee region of Ceylon. — *Ann. Mag. nat. Hist.* (13) 2: 309-320.
KOHN, A. J. 1960b. Spawning behaviour, egg masses and larval development in *Conus* from the Indian Ocean. — *Bull. Bingham oceanogr. Coll.* 17 (4): 1-51.
THIEL, J. 1910. Mollusca: B. Polyplacophora, Gastropoda marina, Bivalvia. In: SCHULTZE, L. *Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Süd-Afrika* 4: 269-270. Jena: Fischer. — *Denkschr. med.-naturw. Ges. Jena* 16: 269-270.

(continued inside back cover)

ANNALS OF THE SOUTH AFRICAN MUSEUM
ANNALE VAN DIE SUID-AFRIKAANSE MUSEUM

Volume 72 Band
December 1976 December
Part 1 Deel



A NEW SPECIES OF *BRADYIDIUS* (COPEPODA,
CALANOIDA) FROM THE MGAZANA ESTUARY,
PONDOLAND, SOUTH AFRICA, AND A REVIEW OF
THE CLOSELY RELATED GENUS *PSEUDOTHARYBIS*

By

JANET M. BRADFORD

Cape Town

Kaapstad

The ANNALS OF THE SOUTH AFRICAN MUSEUM

are issued in parts at irregular intervals as material
becomes available

Obtainable from the South African Museum, P.O. Box 61, Cape Town

Die ANNALE VAN DIE SUID-AFRIKAANSE MUSEUM

word uitgegee in dele op ongereelde tye na beskikbaarheid
van stof

Verkrygbaar van die Suid-Afrikaanse Museum, Posbus 61, Kaapstad

OUT OF PRINT/UIT DRUK

1, 2(1, 3, 5-8), 3(1-2, 4-5, t.-p.i.), 5(1-3, 5, 7-9),
6(1, t.-p.i.), 7(1-4), 8, 9(1-2), 10(1),
11(1-2, 5, 7, t.-p.i.), 15(5), 24(2), 27, 31(1-3), 33

Price of this part/Prys van hierdie deel
R1,70

Trustees of the South African Museum © Trustees van die Suid-Afrikaanse Museum
1976

ISBN 0 949940 98 4

Printed in South Africa by
The Rustica Press, Pty., Ltd.,
Court Road, Wynberg, Cape

In Suid-Afrika gedruk deur
Die Rustica-pers, Edms., Bpk.,
Courtweg, Wynberg, Kaap

A NEW SPECIES OF *BRADYIDIUS* (COPEPODA, CALANOIDA)
FROM THE MGAZANA ESTUARY, PONDOLAND, SOUTH AFRICA,
AND A REVIEW OF THE CLOSELY RELATED GENUS
PSEUDOTHARYBIS

By

JANET M. BRADFORD

*New Zealand Oceanographic Institute, Department of Scientific
and Industrial Research, Wellington*

(With 5 figures and 1 table)

[MS. accepted 28 April 1976]

ABSTRACT

A new species, *Bradyidius hirsutus*, is described. Four species previously attributed to *Bradyidius* are transferred to *Pseudotharybis* which is placed in the Aetideidae.

CONTENTS

	PAGE
Introduction	1
Description of material	2
Review of <i>Pseudotharybis</i>	9
Acknowledgements	9
References	9

INTRODUCTION

The nine previously known species of the aetideid genus *Bradyidius* are *B. angustus* (Tanaka, 1957), *B. arnoldi* Fleminger, 1957, *B. bradyi* (Sars, 1902) (see Matthews (1964) for discussion of priority of specific name *bradyi* over *armatus*), *B. luluae* Grice, 1972, *B. pacificus* (Brodsky, 1950), *B. saanichi* Park, 1960, *B. similis* (Sars, 1902), *B. spinifer* Bradford, 1969a, and *B. tropicus* Wolfenden, 1905. Both sexes are known except for *B. angustus* described only from the male, and *B. tropicus* described only from the female and which has not been morphologically distinguished from *B. bradyi* except that it is smaller and has fewer, coarser teeth on the terminal exopod spine of legs 2-4 (Wolfenden 1905: 1006). At the end of this paper four species, previously attributed to *Bradyidius*, are removed to another genus.

The new species of *Bradyidius* described below was found in plankton samples from the Mgzana Estuary, Pondoland, South Africa (31° 50' S). Although *Bradyidius* is known to be a benthic genus these particular specimens were taken by T. Wooldridge on 1 September 1972 at night just below the surface on an incoming tide at the mouth of the Mgzana Estuary. The depth at this locality is about 3 metres at low tide, salinity 35.0‰ (measured by hydrometer) and temperature 17.8°C.

Dissected specimens of the presently described new species of *Bradyidius* were mounted unstained in Euparal and observed using phase contrast. Descriptions have been made from paratype material.

DESCRIPTION OF MATERIAL

Bradyidius hirsutus sp. nov.

(Figs 1-5)

Holotype

SAM-A13661 in the South African Museum, Cape Town. Adult female in plankton sample from mouth of Mgazana Lagoon, collected by T. Wooldridge, 1 September 1972.

Paratypes

SAM-A13662 in the South African Museum, Cape Town; 1 male whole, 2 males and 2 females dissected, collection data as above.

BM(NH)1975.1124 (1 female), BM(NH)1975.1125 (1 male) in the British Museum (Natural History), London; collection data as above.

P295 in the New Zealand Oceanographic Institute, Wellington, collection; 1 female and 1 male, collection data as above.

Description

Female: Holotype 1.61 mm, paratype specimens 1.59-1.68 mm.

Head and pedigerous segment 1 separate, pedigerous segments 4 and 5 fused. Rostrum bifurcate with slightly divergent points. Posterior metasomal points extend not quite to posterior border of genital segment.

Antenna 1 does not extend quite to posterior metasomal points, of 24 joints, each with annulate setae, joints 8 and 9 fused, joints 18-25 bear very long annulate setae as typical for the genus.

Antenna 2. Endopod joint 1 with 1 seta, joint 2 with 7 outer setae, 7 inner setae (1 small); exopod joint 1 with 1 small seta and joint 2 with 3 setae.

Mandible. Basipod 2 with 1 seta; endopod joint 1 with 1 large and 2 small setae, joint 2 with 10 setae, 1 of them small.

Maxilla 1. Inner lobe 1 with 14 spines and setae, inner lobe 2 with 5 setae, inner lobe 3 and 4 setae, basipod 2 with 5 setae, endopod with at least 13 setae, exopod with 10 setae, outer lobe 1 with 9 setae and outer lobe 2 naked.

Maxilla 2. Lobes 1-5 each with 3 spine-like setae, lobe 5 with 1 of these spines very thick and heavy. Terminal part of limb with 7 setae.

Maxilliped as in other *Bradyidius*.

Leg 1. Basipod 1 with outer edge patch of spines, outer edge spines of exopod joints 1 and 2 not extending beyond base of spine on next joint, outer edge spine on exopod joint 3 longer than its joint.

Leg 2. Basipod 1 with outer edge patch of spines; anterior surface of endopod naked, posterior surface of endopod joint 2 with small hairs covering about half area. Terminal exopod spine with 17 or 21 teeth.




Fig. 1. *Bradyidius hirsutus* sp. nov. Female.

A. Lateral view. B. Dorsal view. C. Rostrum. D. Genital segment, lateral view. E. Urosome, dorsal view.

Leg 3. Basipod 1 with outer edge patch of spines; anterior surface of endopod joints 2 and 3 with small hairs covering slightly smaller area than spines on posterior surface. Terminal exopod spine with 18 or 19 teeth.

Leg 4. Posterior surface of endopod joints 2 and 3 also anterior surface of endopod joint 3 with small hairs. Terminal exopod spine with 20–23 teeth.

Leg 5 absent.

Male: Paratypes 1.36–1.53 mm.

Head and pedigerous segment 1 also pedigerous segments 4 and 5 completely fused. Rostrum bifurcate with slightly diverging points. Posterior metasomal points extend as far as posterior border of urosome segment 1.

Antenna 1 does not extend beyond posterior metasomal points; of 21 or 22 joints, most with annulate setae, 8–10, 12–13 fused as are 20–21 on right side; joints 1–12 with large aesthetes.

Antenna 2 similar to that of female except basipod 1 with row of conspicuous hairs, exopod joint 1 without setae, endopod joint 2 with 2 setae; endopod joint 2 with 6 outer setae.

Mandible blade without teeth, basipod apparently without seta, endopod joint 2 with 10 setae.

Maxilla 1 inner lobes 1-3 without spines or setae, endopod with a few rudimentary setae, exopod and outer lobe 1 with setae.

Maxilla 2 reduced to small lobe without setae.

Maxilliped similar to that of female except basipod 1 has 2 setae distally.

Legs 1-4 similar to those of female except outer edge exopod spines smaller. Terminal spine of legs 2-4 with 20-23, 20-22, and 24-26 teeth respectively.




Fig. 2. *Bradyidius nursutus* sp. nov. Female.

A. Antenna 1. B. Antenna 2. C. Mandible. D. Maxilla 1. E. Maxilla 2. F. Maxilliped.




Fig. 3. *Bradyidius hirsutus* sp. nov. Female.

A. Leg 1, anterior view. B. Leg 2, posterior view. C. Leg 3, anterior view and posterior view of endopod. D. Leg 4, posterior view and anterior view of endopod.

Leg 5 uniramous on each side, hardly extending beyond caudal rami, right leg styliform, of 4 joints; left leg of 5 joints, penultimate joint with inner distal knob bearing a fine spinule, terminal joint with row of small spinules.

Etymology

From the Latin *hirsutus* = hairy, referring to the fact that hairs are to be found on both the anterior and posterior surfaces of legs 3 and 4.

Discussion

All the species of *Bradyidius* are compared in Table 1 in so far as they are known.

Female *Bradyidius hirsutus* may be distinguished primarily from *B. arnoldi*, *B. saanichi* and *B. spinifer*, the three species it most resembles in respect to the length of the posterior metasomal points in that the anterior surfaces of legs 3 and 4 as well as posterior endopod surfaces of legs 2–4 are ornamented with many small hairs. Also the rostral points of *B. hirsutus* are slightly divergent but in *B. arnoldi* not divergent and in *B. saanichi* strongly divergent (exact shape not

TABLE 1
Comparison of species of *Bradydinus*

Species	Length (mm)	FEMALE			FEMALE + MALE		MALE
		Extent of posterior metasomal processes	P ₁ outer edge spine of Re ₁	Anterior-posterior surface spines Ri P ₂₋₄	Rostrum	P ₅	
<i>B. angustus</i>	♀ unknown ♂ 1,37	—	—	Posterior	Hardly divergent	Uniramous: distal border right Re ₂ extends just beyond distal border left B ₂	
<i>B. arnoldi</i>	♀ 1,64 ♂ 1,19-1,23	Almost to posterior border genital segment	Very small	Posterior	Not divergent	Biramous	
<i>B. bradyi</i>	♀ 2,65 ♂ 2,20	Posterior border genital segment	Reaches base next spine	Posterior	Slightly divergent	Uniramous: right leg much shorter than left	
<i>B. luluae</i>	♀ 3,12 ♂ 3,00-3,25	Just beyond posterior border genital segment	Extends half way along next spine	Posterior (♂)	Strongly divergent	Uniramous: distal border right Re ₂ extends to middle left Re ₁	
<i>B. pacificus</i>	♀ 4,4 ♂ 3,3	Terminal one-third genital segment	Reaches base next spine	None	Not divergent	Biramous	
<i>B. saanichi</i>	♀ 2,33-2,56 ♂ 2,01-2,24	Almost to posterior border genital segment	Reaches base next spine	None	Strongly divergent	Biramous	
<i>B. similis</i>	♀ 3,0 ♂ 2,4	Hardly beyond middle genital segment	As in <i>B. bradyi</i>	None	—	Biramous	
<i>B. spinifer</i>	♀ 2,74 ♂ 1,96	Almost to posterior border genital segment	Extends over half way along next spine	Posterior	—	Uniramous: distal border right Re ₂ extends to middle left Re ₁	
<i>B. tropicus</i>	♀ 1,2 ♂ unknown	—	—	—	—	—	
<i>B. hirsutus</i> sp. nov.	♀ 1,59-1,68 ♂ 1,36-1,53	Almost to posterior border genital segment	Does not reach base next spine	Posterior and anterior	Slightly divergent	Uniramous: distal border right Re ₂ extends just beyond distal border left B ₂	

— information not known; Re_{1,2} = exopod joint 1, 2; P₁₋₅ = legs 1-5; Ri = endopod; B₂ = basipod 2.

— information not known; Re_{1,2} = exopod joint 1, 2; P₁₋₅ = legs 1-5; Ri = endopod; B₂ = basipod 2.

known for *B. spinifer*) and the outer edge spine of leg 1 exopod joint 1 does not reach the base of the next spine in *B. hirsutus*, is very small in *B. arnoldi*, reaches the base of the next spine in *B. saanichi* and extends half-way along the next spine in *B. spinifer*.

The male of *B. hirsutus* is most like that of *B. angustus* even to the proportions of leg 5 but differs from it chiefly in the ornamentation on the anterior and posterior surfaces of legs 3 and 4. Also exopod joint 1 of antenna 2 has no setae in *B. hirsutus* whereas it appears to have a seta in Tanaka's (1957) illustration and the rostrum of *B. hirsutus* has a much wider notch at the base of the points than in *B. angustus*.

It appears the setation of the female maxilla 1 may be conservative. Re-examination of the holotype of *B. spinifer* revealed that the basipod 2 in fact has 5 setae, not 4 as described by Bradford (1969a) so *B. spinifer* does not differ from *B. lulae* in this respect.




Fig. 4. *Bradyidius hirsutus* sp. nov. Male.

A. Lateral view. B. Dorsal view. C. Rostrum. D. Urosome, dorsal view.




Fig. 5. *Bradyidius hirsutus* sp. nov. Male.

A. Left antenna 1. B. Antenna 2. C. Mandible. D. Maxilla 1. E. Maxilliped and rudimentary maxilla 2. F. Leg 1, anterior view. G. Leg 2, anterior view and posterior view of endopod. H. Leg 3, anterior view and posterior view of endopod. I. Leg 4, anterior view and posterior view of endopod. J. Leg 5.

REVIEW OF *PSEUDOTHARYBIS*

Females of three species attributed to *Bradyidius* by Bradford (1969b), *B. robustus*, *B. brevispinus* and a stage V *B. spinibasis* (erroneously referred to as *B. spinatus* in the text (Bradford 1969a: 484)) have subsequently been recognized by the author as fitting T. Scott's (1909a) brief description of *Pseudotharybis* because of the unique form of the female fifth legs which hardly varies from one species to another. Although T. Scott (1909a) implied *Pseudotharybis* was a tharybid, lack of sensory appendages terminally on maxilla 2 precluded that conclusion. This genus is here placed in the family Aetideidae, closely related to *Bradyidius* and *Aetideopsis*, and is one of the few genera in the family to have a female fifth leg.

The male of *B. robustus* Bradford, 1969a along with *B. dentatus* Bradford, 1969a indicates the form of male *Pseudotharybis* which hardly differs from that of *Bradyidius* or *Aetideopsis* except that the posterior metasomal points are represented in *Pseudotharybis* by a small posterodorsal tooth and in the other two genera by a posterior spine. Thus the genus *Pseudotharybis* now contains the following species: *P. zetlandicus* T. Scott, 1909a, *P. brevispinus* (Bradford, 1969a), *P. dentatus* (Bradford, 1969a), *P. robustus* (Bradford, 1969a), and *P. spinibasis* (Bradford, 1969a). There is also a possibility that *Aetideopsis magna* Grice & Hulsemann, 1970 belongs to this genus. Pending a closer examination of the genera *Aetideopsis*, *Bradyidius* and *Pseudotharybis* the exact limits of each genus, especially with reference to the males, remains uncertain.

T. Scott's (1909b) *P. dubius* must be removed from *Pseudotharybis* as there appear to be important differences from all other species in the genus. For example maxilla 1 inner and outer lobe 1 are well developed but the remaining parts are reduced in size and have a smaller number of setae, maxilliped basipod 1 has a strong spiniform seta on its distal border and leg 5 is of a different shape.

ACKNOWLEDGEMENTS

I am indebted to Mr T. Wooldridge of the Port Elizabeth Museum for bringing the new species of *Bradyidius* to my attention.

REFERENCES

- BRADFORD, J. M. 1969a. New species of *Aetideopsis* Sars and *Bradyidius* Giesbrecht (Copepoda: Calanoida) from the Southern Hemisphere. — *N.Z. Jl mar. freshw. Res.* 3: 73–97.
- BRADFORD, J. M. 1969b. New genera and species of benthic calanoid copepods from the New Zealand slope. *N.Z. Jl mar. freshw. Res.* 3: 473–505.
- BRODSKY, K. A. 1950. Calanoida of the far eastern seas and polar basin of the U.S.S.R.—*Opred. Faune SSSR* 35: 1–442. (Translation: Israel Program for Scientific Translation, Jerusalem, 1967).
- FLEMINGER, A. 1957. New calanoid copepods of the families Aetideidae, Euchaetidae, and Stephidae from the Gulf of Mexico.—*Fishery Bull. Fish Wildl. Serv. U.S.* 57: 355–63.
- GRICE, G. D. 1972. The existence of a bottom-living calanoid copepod fauna in deep water with descriptions of five new species.—*Crustaceana* 23: 219–42.

- GRICE, G. D. & HULSEMAN, K. 1970. New species of bottom-living calanoid copepods collected in deep water by the DSRV Alvin. — *Bull. Mus. comp. Zool. Harv.* **139**: 185–227.
- MATTHEWS, J. B. L. 1964. On the biology of some bottom-living copepods (Aetideidae and Phaennidae) from western Norway. — *Sarsia* **16**: 1–46.
- PARK, T. 1960. A new species of *Bradyidius* (Copepoda: Calanoida) from the Pacific coast of North America. — *J. Fish Res. Bd Canada* **23**: 805–11.
- SARS, G. O. 1902, 1903. Copepoda. In: *An account of the Crustacea of Norway* **4**. Bergen: Bergen Museum.
- SCOTT, T. 1909a. On some new and rare Entomostraca from the Scottish Seas. — *Ann. Mag. nat. Hist.* (8) **3**: 122–30.
- SCOTT, T. 1909b. On new and rare Crustacea from Scottish waters. — *Ann. Mag. nat. Hist.* (8) **4**: 31–6.
- TANAKA, O. 1957. The pelagic copepods of the Izu region, middle Japan. Systematic account III. Family Aetideidae (Part I). — *Publs Seto mar. biol. Lab.* **6**: 31–68.
- WOLFENDEN, R. N. 1905. Notes on the collection of Copepoda. In: GARDINER, J. S. (ed.) *The Fauna and Geography of the Maldiv and Laccadive Archipelagoes* **2**: suppl. 1: 989–1040. Cambridge: University Press; London: Clay.

6. SYSTEMATIC papers must conform with the *International code of zoological nomenclature* (particularly Articles 22 and 51).

Names of new taxa, combinations, synonyms, etc., when used for the first time, must be followed by the appropriate Latin (not English) abbreviation, e.g. gen. nov., sp. nov., comb. nov., syn. nov., etc.

An author's name when cited must follow the name of the taxon without intervening punctuation and not be abbreviated; if the year is added, a comma must separate author's name and year. The author's name (and date, if cited) must be placed in parentheses if a species or subspecies is transferred from its original genus. The name of a subsequent user of a scientific name must be separated from the scientific name by a colon.

Synonymy arrangement should be according to chronology of names, i.e. all published scientific names by which the species previously has been designated are listed in chronological order, with all references to that name following in chronological order, e.g.:

Family Nuculanidae

Nuculana (Lembulus) bicuspidata (Gould, 1845)

Figs 14–15A

Nucula (Leda) bicuspidata Gould, 1845: 37.

Leda plicifera A. Adams, 1856: 50.

Laeda bicuspidata Hanley, 1859: 118, pl. 228 (fig. 73). Sowerby, 1871: pl. 2 (figs 8a–b).

Nucula largillierti Philippi, 1861: 87.

Leda bicuspidata: Nicklès, 1950: 163, fig. 301; 1955: 110. Barnard, 1964: 234, figs 8–9.

Note punctuation in the above example:

comma separates author's name and year

semicolon separates more than one reference by the same author

full stop separates references by different authors

figures of plates are enclosed in parentheses to distinguish them from text-figures

dash, not comma, separates consecutive numbers

Synonymy arrangement according to chronology of bibliographic references, whereby the year is placed in front of each entry, and the synonym repeated in full for each entry, is not acceptable.

In describing new species, one specimen must be designated as the holotype; other specimens mentioned in the original description are to be designated paratypes; additional material not regarded as paratypes should be listed separately. The complete data (registration number, depository, description of specimen, locality, collector, date) of the holotype and paratypes must be recorded, e.g.:

Holotype

SAM-A13535 in the South African Museum, Cape Town. Adult female from mid-tide region, King's Beach, Port Elizabeth (33.51S, 25.39E), collected by A. Smith, 15 January 1973.

Note standard form of writing South African Museum registration numbers, date and geographical positions.

7. SPECIAL HOUSE RULES

Capital initial letters

- (a) The Figures, Maps and Tables of the paper when referred to in the text
e.g. '... the Figure depicting *C. namacolus* ...'; '... in *C. namacolus* (Fig. 10) ...'
- (b) The prefixes of prefixed surnames in all languages, when used in the text, if not preceded by initials or full names
e.g. Du Toit but A. L. du Toit; Von Huene but F. von Huene
- (c) Scientific names, but not their vernacular derivatives
e.g. Therocephalia, but therocephalian

Punctuation should be loose, omitting all not strictly necessary

Reference to the author should be expressed in the third person

Roman numerals should be converted to arabic, except when forming part of the title of a book or article, such as

'Revision of the Crustacea. Part VIII. The Amphipoda.'

Specific name must not stand alone, but be preceded by the generic name or its abbreviation to initial capital letter, provided the same generic name is used consecutively.

Name of new genus or species is not to be included in the title: it should be included in the abstract, counter to Recommendation 23 of the Code, to meet the requirements of Biological Abstracts.



JANET M. BRADFORD

A NEW SPECIES OF *BRADYIDIUS* (COPEPODA,
CALANOIDA) FROM THE MGAZANA ESTUARY,
PONDOLAND, SOUTH AFRICA, AND A
REVIEW OF THE CLOSELY RELATED GENUS
PSEUDOTHARYBIS